




- Complete this sentence: The transformation that maps shape A onto **shape B** is **translate** ..... squares to the left.



- Translate shape F four squares to the left.



- Translate shape V two squares to the left and one square up.




4.

4. not written yet

5. ABCDE is an irregular pentagon.

Write down the vector  $\vec{AB}$


$$\vec{AB} = \begin{pmatrix} \dots \\ \dots \end{pmatrix}$$



6.

6. not written yet


7. Here are some vectors.



Write down the column vectors for

(i)  $\mathbf{a} = \begin{pmatrix} \phantom{\dots} \\ \phantom{\dots} \end{pmatrix}$     (ii)  $-\mathbf{a} = \begin{pmatrix} \phantom{\dots} \\ \phantom{\dots} \end{pmatrix}$     (iii)  $2\mathbf{a} = \begin{pmatrix} \phantom{\dots} \\ \phantom{\dots} \end{pmatrix}$     (iv)  $\vec{SR} = \begin{pmatrix} \phantom{\dots} \\ \phantom{\dots} \end{pmatrix}$

8. On the grid below, translate trapezium V by the vector  $\begin{bmatrix} 2 \\ -1 \end{bmatrix}$  and label it W




9.  $\mathbf{p} = \begin{pmatrix} 5 \\ -3 \end{pmatrix}$        $\mathbf{q} = \begin{pmatrix} -2 \\ 1 \end{pmatrix}$

Complete these column vectors

(i)  $2\mathbf{p} = \begin{pmatrix} \dots \\ \dots \end{pmatrix}$       (ii)  $3\mathbf{q} = \begin{pmatrix} \dots \\ \dots \end{pmatrix}$

10. (a) The diagram below shows 12 congruent parallelograms.



$\vec{OA} = \mathbf{a}$  {Handwriting **bold** is hard to do so mathematicians write a instead of **a**}


$\vec{OD} = \mathbf{d}$  {and write d instead of **d**}

Find in terms of **a** and **d** the vectors

(i)  $\vec{AC} = \dots\dots\dots$

(ii)  $\vec{GU} = \dots\dots\dots$

(b) The diagram below shows parallelogram ABCD



The diagonals of the parallelogram intersect at O

$\vec{OA} = \mathbf{a}$  {Handwriting **bold** is hard to do so mathematicians write a instead of **a**}

$\vec{OB} = \mathbf{b}$  {and write b instead of **b**}

Write an expression, in terms of **a** and **b** for

(i)  $\vec{CO} = \dots\dots\dots$

(ii)  $\vec{DB} = \dots\dots\dots$

$$11. \quad \mathbf{a} = \begin{pmatrix} -2 \\ -1 \end{pmatrix} \quad \mathbf{b} = \begin{pmatrix} 3 \\ -4 \end{pmatrix}$$

Work out  $\mathbf{a} + \mathbf{b}$  as a column vector.

$$\begin{pmatrix} \dots \\ \dots \end{pmatrix}$$

$$12. \quad \mathbf{a} = \begin{pmatrix} 2 \\ -1 \end{pmatrix} \quad \mathbf{b} = \begin{pmatrix} -2 \\ 3 \end{pmatrix}$$

Work out  $3\mathbf{a} + 2\mathbf{b}$  as a column vector.

$$\begin{pmatrix} \dots \\ \dots \end{pmatrix}$$